

# DETERMINATION OF LEAD CONTENT IN WHITENING CREAM USING LASER INDUCED BREAKDOWN SPECTROSCOPY ANALYSIS

WAN NORSYUHADA BT WAN MOHD SHUKRI

A thesis submitted in fulfillment of the  
requirements for the award of the degree of  
Master of Science (Physics)

Faculty of Science  
Universiti Teknologi Malaysia

SEPTEMBER 2014

## DEDICATION

*“I humbly thank Allah Almighty, the Merciful and the Beneficent, who gave me health, thoughts and co-operative people to enable me achieve this goal.”*

***To My Dearest Husband***

*Mohd Azahar b Che Abdullah*

*Thank you so much for your love and motivation*

***To My Beloved Parents,***

*W. Mohd Shukri b W. Ibrahim & Huzian bt Husin*

*Thank you very much for your love and supports*

***To My Lovely Siblings***

*W. Muhammad Syafiq b W. Mohd Shukri*

*W. Syazwan b W. Mohd Shukri*

*W. Nursyamimi bt W. Mohd Shukri*

***To My Respected and Beloved Supervisor***

*Prof. Dr. Noriah Ibrahim*

*Thank you very much for your attention and guidance*

***And all my friends...***

*Especially Nurul Hida and Saleha....*

## ACKNOWLEDGEMENT

I humbly thank Allah Almighty, the Merciful and the Beneficent, who gave me health, thoughts and co-operative people to enable me achieve this goal. It is a pleasure to thank the many people who made this thesis possible.

I wish to thank, first and foremost to my great thesis supervisor, Prof. Dr. Noriah Bidin. Throughout my thesis-writing periods she provided encouragement, sound advice, good teaching, good company, and lots of good ideas. I would have been lost without her.

Thanks to Mrs Shima and Mrs Sarah for their cooperation during my research process especially in laboratory equipment's. Thanks to all lab mates, especially Saleha, Hida, Radhiana, Daeng, Aiza, Mudzir, Shafiq and Mustafa for being helpful during my research study in the lab.

I am most grateful to Mr Subri as the Institute Bioproduct Development's Assistant for being helpful me to get the research results using ICP-MS.

Lastly, I would like to thank to UTM for giving me an opportunity to get more experience in my master degree.

## ABSTRACT

Laser Induced Breakdown Spectroscopy (LIBS) technique was applied to determine the concentration of heavy metal such as Pb in five different whitening creams available from retail shop and open market in Johor, Malaysia. The whitening cream in the form of an emulsion was painted on the glass slide and then cryogenically frozen in refrigerator. The harden sample of whitening creams were irradiated with light Q-switch Nd:YAG laser at five different positions on the same sample. The spectral lines of six whitening creams were identified using National Institute of Standards and Technology (NIST) Database. Finally, Pb was detected in all the studied samples and their concentration in the range of 1.78 ppm – 4.64 ppm with limit of detection which is  $\sim 1$  ppm. The LIBS results were validated by Inductive Couple Plasma Mass Spectroscopy (ICP-MS) technique with percentage difference in the range of 2.73% – 7.60%. The detection of Pb in the whitening creams is under permissible limit as set by Malaysian standard. However, the limitation is over as set by WHO which is 0.05 ppm for daily used. Thus, it is better to take precaution because continuously using creams may lead to accumulative toxicity in the body beyond the acceptable limit. Thus, the high sensitivity, portable and *in-situ* system allow LIBS to be an important technique in cosmetic industries.

## ABSTRAK

Teknik spektroskopi leraian aruhan laser (LIBS) telah digunakan untuk menentukan kepekatan logam berat Pb di dalam enam jenis krim pemutih yang berbeza yang boleh didapati dari kedai runcit dan pasaraya di Malaysia. Krim pemutih adalah dalam bentuk emulsi telah dicat pada kepingan kaca dan kemudian dibekukan dalam peti sejuk. Sampel krim yang telah dikeraskan kemudian diradiasikan dengan cahaya laser suis-Q Nd : YAG di lima kedudukan yang berbeza pada tempat sasaran yang sama. Spektrum bagi enam jenis krim pemutih telah dikenal pasti dengan menggunakan data dari Institut Piawaian dan Teknologi Kebangsaan (NIST). Akhir sekali, Pb dikesan dalam semua sampel yang telah dikaji dengan kepekatan dalam lingkungan 1.78 pm – 4.64 ppm dengan had pengesanan ialah ~ 1 ppm. Keputusan LIBS telah disahkan oleh teknik plasma berganding aruhan spektrometri jisim (ICP-MS) dengan perbezaan peratus dalam lingkungan 2.73% – 7.60%. Pb yang dikesan dalam semua krim pemutih adalah di bawah had yang dibenarkan seperti yang telah ditetapkan oleh Piawaian Malaysia. Walau bagaimanapun, had ini adalah berlebihan seperti yang telah ditetapkan oleh WHO iaitu 0.05 ppm untuk kegunaan harian. Oleh itu, adalah lebih baik untuk mengambil langkah berjaga-jaga kerana penggunaan krim ini yang berterusan boleh menyebabkan pengumpulan toksik di dalam badan melebihi had yang boleh diterima. Oleh itu, kepekaan yang tinggi, mudah alih dan sistem *in-situ* membolehkan LIBS sebagai teknik penting dalam industri kosmetik.